Preparation for Colonoscopy: MIRALAX

Location: Endoscopy Center 2nd Floor

1140 W La Veta Orange CA 92868

Appointment Date:	Arrival Time:	

What you need:

- 1) Dulcolax tablets 4 tablets (called in to your pharmacy).
- 2) Miralax 238-gram bottle (found over the counter at any pharmacy).
- 3) 64ozs of a clear liquid as in Gatorade, Crystal Lite, water. You may have: Yellow, Blue, Green or Clear. You cannot have RED, PURPLE or ORANGE colors

<u>Five days prior to your procedure:</u> If you are on Coumadin or Plavix, stop taking these medications. Continue to take all other medications as usual including the day of the procedure. If you are having an EGD the last oral intake has to be 4 hours before the procedure time.

<u>Four days prior to your procedure:</u> Stop Iron and High Fiber foods, this means wheat bread, oat meal, bran, fruit, vegetables, nuts.(This also includes supplements) **You may have** proteins, starches and dairy products.

Two days prior to the procedure: Have breakfast, lunch and a **light dinner**, then begin a clear liquid diet (Broth, Jello, juices, popsicles, tea, sports drinks, clear sodas) they may be blue, yellow or green. You'll continue this diet up until 4 hours prior to your procedure time. NO RED, ORANGE OR PURPLE LIQUIDS ALLOWED.

IF YOU ARE A DIABETIC DO NOT DRINK ANY DIET LIQUIDS IT MAY CAUSE YOUR SUGAR TO DROP.

One day prior to the procedure: You may start your preparation as early as you would like to but no later than 4pm. First take 2 Dulcolax tablets. One hour later mix the entire bottle of Miralax with 64oz of clear liquid, shake until Miralax is dissolved. Slowly drink an 8oz glass of the mixture every 15 to 20 minutes until 32oz are consumed. You may refrigerate the difference. One hour later take the last 2 Dulcolax tablets. Wait four hours then complete the second dose of Miralax the same way you took the first one.

OR

You can finish the last 32ozs of Miralax the day of your procedure the same way you drank it the day before but you need to finish it 4 hours before your arrival time.

IF YOUR ARE STILL SEEING SOLID STOOL AT THE END OF THE PREP YOU NEED TO TAKE AN EXTRA 1/2 DOSE OF THE MIRALAX.

Monitor the urine color it should be a clear yellow. If it is dark you need to increase your fluid intake. You may use Dessitin or Vaseline cream to the anal area to protect the skin. YOU MAY NOT HAVE ANYTHING TO EAT OR DRINK 4 HOURS BEFORE YOUR PROCEDURE TIME.

Important Notes:

You CANNOT drive after the colonoscopy till the following day.

You have 72 business hours to reschedule, if you do not show up to your appointment without prior cancellation there will be \$100.00 cancellations fee NO EXCEPTIONS.

IF A BIOPSY WAS TAKEN PLEASE CALL OUR OFFICE 5-10 DAYS AFTER YOUR PROCEDURE FOR YOUR RESULTS.